

Preprint 98 – 034

**A Combinatorial Algorithm
for the Minimum $(2, r)$ -Metric Problem
and Some Generalizations**

by

Alexander V. Karzanov

Mitgeteilt von W. Deuber
29.4.1998

A combinatorial algorithm for the minimum $(2, r)$ -metric problem and some generalizations

Alexander V. KARZANOV *

Abstract. Let $G = (V, E)$ be a graph with nonnegative integer capacities $c(e)$ of the edges $e \in E$, and let μ be a metric that establishes distances on the pairs of elements of a subset $T \subseteq V$. In the *minimum 0-extension problem* (*), one is asked for finding a metric m on V such that m coincides with μ within T , each $x \in V$ is at zero distance from some $t \in T$, and the value $\sum(c(e)m(e) : e \in E)$ is as small as possible. When $T = \{s, t\}$ and $\mu(s, t) = 1$, this turns into the classical minimum (undirected) cut problem. When μ is the path metric of the complete bipartite graph $K_{2,r}$, (*) is specified to be the *minimum $(2, r)$ -metric problem*. It is known that such a problem can be solved in strongly polynomial time by use of the ellipsoid method.

We develop a polynomial time algorithm for the minimum $(2, r)$ -metric problem, using only “purely combinatorial” means. The algorithm simultaneously solves a certain associated integer multiflow problem. We then apply this algorithm to solve (*) for a wider class of metrics μ , give other results and raise open questions.

Key words: finite metric, cut, metric extension, multicommodity flow.

* On leave from the Institute for System Analysis, 9, Prospect 60 Let Oktyabrya, 117312 Moscow, Russia; email: sasha@cs.isa.ac.ru.

1. Introduction

By a *metric* on a set S we mean a function $d : S \times S \rightarrow \mathbf{R}_+$ that establishes *distances* on the pairs of elements of S satisfying (i) $d(x, x) = 0$, (ii) $d(x, y) = d(y, x)$, and (iii) $d(x, y) + d(y, z) \geq d(x, z)$, for all $x, y, z \in S$. We usually write $d(xy)$ in place of $d(x, y)$ and allow $d(xy) = 0$ for some $x \neq y$. A special case of metrics is the *path metric* d^Γ of a connected graph $\Gamma = (S, W)$, i.e., $d^\Gamma(xy)$ is the minimum number of edges of a path in Γ connecting x and y .

We consider an undirected graph $G = (V, E)$ whose edges $e \in E$ have nonnegative integer *capacities* $c(e)$, and a subset $T \subseteq V$ of nodes called *terminals*. Let μ be a metric on T . A metric m on V is said to be an *extension* of μ to V if the submetric of m on T is just μ (i.e., m and μ coincide within T), and to be a *0-extension* if, in addition, for each $x \in V$, there is $t \in T$ such that $m(tx) = 0$. If μ is a *positive* metric, i.e., $\mu(st) > 0$ for any $s \neq t$, then each 0-extension m of μ to V one-to-one corresponds to a *T-partition*, i.e., a partition of V into $|T|$ subsets $\{X_t : t \in T\}$ where each X_t contains exactly one terminal, namely, t . This correspondence is defined by setting $X_t = \{x \in V : m(tx) = 0\}$.

A path in G whose ends are different elements of T is called a *T-path*. A (c -admissible) *multiflow* f consists of T -paths P_1, \dots, P_k in G along with nonnegative real weights $\lambda_1 = \lambda(P_1), \dots, \lambda_k = \lambda(P_k)$ satisfying the capacity constraint

$$f^e := \sum (\lambda_i : P_i \text{ contains } e) \leq c(e) \quad \text{for all } e \in E.$$

Define $\langle \mu, f \rangle$, the μ -value of f , to be $\sum (\mu(u_i v_i) \lambda_i : i = 1, \dots, k, u_i \text{ and } v_i \text{ are the ends of } P_i)$. If all λ_i 's are integers, f is called an *integer* multiflow. Consider the following four problems:

- (1.1) Find a 0-extension m of μ to V with $c \cdot m := \sum (c(e)m(e) : e \in E)$ as small as possible;
- (1.2) Find an extension m of μ to V with $c \cdot m$ as small as possible;
- (1.3) Find an integer multiflow f whose μ -value is as large as possible;
- (1.4) Find a multiflow f whose μ -value is as large as possible.

(Problem (1.1) is also known as the multifacility location problem, cf. [TFL].) Let $\tau = \tau(G, c, \mu)$ and $\tau^* = \tau^*(G, c, \mu)$ denote the minimum $c \cdot m$ in (1.1) and (1.2), respectively, and let $\nu = \nu(G, c, \mu)$ and $\nu^* = \nu^*(G, c, \mu)$ denote the maximum μ -values in (1.3) and (1.4), respectively. For given G, c, μ , an extension m with $c \cdot m = \tau^*$ is called *minimum* and a multiflow f with $\langle \mu, f \rangle = \nu^*$ is called *maximum*, and similarly for 0-extensions and integer multiflows. Since (1.1) is a strengthening of (1.2), and (1.3)

is a strengthening of (1.4), we have $\tau \geq \tau^*$ and $\nu \leq \nu^*$. In their turn, the relaxations (1.2) and (1.4) are, in fact, mutually dual linear programs (see, e.g., [KL]), whence $\tau^* = \nu^*$. Thus, we have the following relations:

$$(1.5) \quad \tau \geq \tau^* = \nu^* \geq \nu.$$

Each of the two inequalities here may be strict. The simplest case with equality throughout in (1.5) arises when $\mu = d^{K_2}$ (K_p is the complete graph with p nodes). In this case any 0-extension is a cut metric, an optimal 0-extension corresponds to a minimum cut “separating” the pair of terminals, and $\tau = \nu$ holds by the classical max flow min cut theorem [FF]. On the other hand, if $\mu = d^{K_p}$ with $p \geq 3$, then $\tau = \tau^*$ needs not hold; in this case (1.1) turns into the *multiterminal cut problem*, which is known to be strongly NP-hard even if $p = 3$ [Da].

Following [KM], a metric μ on T is called *minimizable* if $\tau(G, c, \mu) = \tau^*(G, c, \mu)$ holds for any graph $G = (V, E)$ with $X \supseteq T$ and capacities $c : E \rightarrow \mathbb{Z}_+$. For such a μ , problem (1.1) can be solved in strongly polynomial time by use of the ellipsoid method, taking into account that (1.2) is, in fact, a linear program whose constraint matrix has a polynomial size in $|V|, |E|$. The class of minimizable metrics is rather large; in particular, it includes $\mu = d^{K_{2,r}}$ for an arbitrary r , where $K_{p,q}$ is the complete bipartite graph whose parts (the maximal stable sets) consist of p and q nodes. For such a μ , a 0-extension is called a $(2, r)$ -metric, and (1.1) can be specified as:

$$(1.6) \quad \text{Given } G, T, c \text{ and a partition of } T \text{ into two subsets } A = \{s_1, s_2\} \text{ and } B = \{t_1, \dots, t_r\}, \text{ find a } T\text{-partition } \{S_1, S_2, T_1, \dots, T_r\} \text{ of } V \text{ with } s_i \in S_i \text{ and } t_j \in T_j \text{ that minimizes } \sum (c(S_i, T_j) : i = 1, 2, j = 1, \dots, r) + 2c(S_1, S_2) + 2 \sum (c(T_i, T_j) : 1 \leq i < j \leq r),$$

where for $X, Y \subseteq V$, $c(X, Y)$ denotes the total capacity of edges with one end in X and the other in Y . Although the second inequality in (1.5) may be strict for $\mu = d^{K_{2,r}}$ too, it holds with equality in the following important case. We say that an (integer) capacity function c on E is *inner Eulerian* if $c(X, V - X)$ is even for each $X \subseteq V - T$.

Theorem 1.1 [KM]. *If c is inner Eulerian and $\mu = d^{K_{2,r}}$, then $\nu = \nu^*$ and, therefore, $\nu = \tau$.*

The main goal of this paper is to give a “purely combinatorial” algorithm which finds a minimum $(2, r)$ -metric and finds a maximum integer multiflow when c is inner Eulerian. The algorithm we develop runs in time polynomial in $|V|, |E|$ and linear in $\log \|c\|$, where $\|c\| = \max\{c(e) : e \in E\}$.

In fact, the algorithm focuses on construction of a maximum integer multiflow, whereas a minimum $(2, r)$ -metric is obtained as a by-product. It involves three ingredients: (i) a capacity scaling method, (ii) an integer augmentation procedure, and (iii)

a maximality check-up procedure. These occur in the high, middle and low levels of the algorithm, respectively. The capacity scaling method reduces the whole problem to about $\log \|c\|$ similar problems each of which deals with a truncated capacity function c' and finds a maximum integer multiflow for it, starting with a nearly optimal integer multiflow f' whose μ -value is only $O(|E|)$ below $\tau(G, c', \mu)$. Therefore, at most $O(|E|)$ integer augmentations of the μ -value are sufficient to transform f' into a maximum integer multiflow for c' . The integer augmentation procedure is somewhat more complicated than that based on standard augmenting path techniques in maximum flow algorithms. It involves a vertex splitting method and relies on possibility to decide whether a given (fractional) multiflow f'' for a capacity function c'' on E is maximum or not. Our maximality check-up procedure solves the latter problem in strongly polynomial time and in a combinatorial fashion.

Next we consider a more general case. A complete characterization of the class of graphs whose path metric is minimizable is given in the following theorem.

Theorem 1.2 [K98]. *For a graph H , the metric d^H is minimizable if and only if H is bipartite, orientable and contains no isometric k -circuit with $k \geq 6$.*

Here a subgraph (or circuit) $H' = (T', U')$ of H is called *isometric* if $d^{H'}$ is a submetric of d^H ; a *k -circuit* is a (simple) circuit with k nodes; and H is called *orientable* if its edges can be oriented so that for any 4-circuit $C = (v_0, e_1, v_1, \dots, e_4, v_4 = v_0)$, the orientations of the opposite edges e_1 and e_3 are different along the circuit (i.e., if e_1 is oriented from v_0 to v_1 say, then e_3 is oriented from v_3 to v_2), and similarly for e_2 and e_4 ; a feasible orientation for C is shown in Fig. 1a. For example, the graph $K_{p,r}$ is orientable if and only if $\min\{p, r\} \leq 2$. A graph H as in Theorem 1.2 is called a *frame* (hereinafter we assume, w.l.o.g., that H has no parallel edges or loops).




Fig. 1

(a)

(b)

We say that edges e, e' of a graph $H = (T, U)$ are *dependent* if there is a sequence $e = e_0, e_1, \dots, e_k = e'$ where each two consecutive e_i, e_{i+1} are opposite edges in some 4-circuit of H . This relation is symmetric and transitive, and a maximal set of dependent edges is called an *orbit* of H . In particular, the edges of each subgraph $K_{2,r}$ in H belong to the same orbit when $r \geq 3$. For a subset $Z \subseteq U$, let H/Z denote the graph obtained from H by contracting each edge in Z .

Definition. A frame $H = (T, U)$ with the orbits Q_1, \dots, Q_k is called *sparse* if for each i , the graph H_i obtained from $H/(U - Q_i)$ by identifying the parallel edges and deleting the loops (if any) is $K_{p,r}$ with $\min\{p, r\} \leq 2$.

Remark. One can show that a frame is sparse if and only if each orbit in it contains (the edge set of) at most one inclusion maximal subgraph $K_{2,r}$ with $r \geq 3$.

Figure 1b illustrates a sparse frame with four orbits. As mentioned above, for any fixed minimizable metric μ , (1.1) is solvable in strongly polynomial time by use of the ellipsoid method. We show that for any sparse frame H , (1.1) with $\mu = d^H$ is reduced to $O(|T|)$ minimum cut or minimum $(2, r)$ -metric computations. This reduction and our minimum $(2, r)$ -metric algorithm provide a “purely combinatorial” polynomial time algorithm for such a μ .

This paper is organized as follows. The algorithm for the minimum $(2, r)$ -metric problem and its corresponding integer multiflow problem in the inner Eulerian case is described throughout Sections 2–4. Section 5 is devoted to the generalization for sparse frames. The concluding Section 6 contains some remarks and raises open questions.

In what follows, for $X \subseteq V$, $\delta(X) = \delta^G(X)$ denotes the set of edges of G with one end in X and the other in $V - X$ (a *cut* in G), and $c(\delta(X))$ stands for $c(X, V - X)$. For a multiflow f in a network (G, c, T) (as above) and a pair $\{u, v\} \subseteq T$, f_{uv} denotes the part of f concerning the paths with the ends u and v (the *flow* in f between u and v); the total weight of these paths is called the *value* of f_{uv} and denoted by $|f_{uv}|$. Hence, the μ -value of f is $\sum(\mu(uv)|f_{uv}| : \{u, v\} \subseteq T)$. For a path P in G , χ^P denotes its incidence vector in \mathbf{R}^E , i.e., $\chi^P(e)$ is the number of occurrences of an edge e in P .

In conclusion of this section we point out one application of the minimum $(2, r)$ -metric problem. In the *multiflow demand problem*, one is given a set D of pairs uv of terminals and *demands* $d(uv) \in \mathbf{Z}_+$ on these pairs, and is asked to find a multiflow f in (G, c, T) satisfying $|f_{uv}| = d(uv)$ for all $uv \in D$. By a five-terminus flow theorem [K87], when $|T| = 5$, the demand problem has a solution if and only if the cut condition

$$(1.7) \quad c(\delta(X)) \geq \sum(d(uv) : uv \in D, |\{u, v\} \cap X| = 1) =: d_X$$

holds for each $X \subset V$, and the $(2,3)$ -metric condition

$$(1.8) \quad c \cdot m \geq \sum(d(uv)m(uv) : uv \in D)$$

holds for each $(2,3)$ -metric m on V (concerning all possible $H = (T, U) \simeq K_{2,3}$). Also [K87] shows that if (c, d) satisfies the parity condition

$$(1.9) \quad c(\delta(X)) + d_X \equiv 0 \pmod{2} \quad \text{for all } X \subseteq V,$$

then the problem has an integer solution f provided that it has a solution at all; moreover, to find such an f takes $O(|V|^3)$ minimum cut and minimum $(2,3)$ -metric

computations. This implies that f can be found in polynomial time by a “purely combinatorial” algorithm.

2. Checking the maximality

Let $f = (P_1, \dots, P_k, \lambda_1, \dots, \lambda_k)$ be a multiflow for G, T, c , and let $\mu = d^H$, where $H = (T, U)$ is $K_{2,r}$. We assume that $\lambda_1, \dots, \lambda_k > 0$. The algorithm described in this section decides whether the given f is maximum, and if so, finds a minimum $(2, r)$ -metric. We apply a “flow expansion” approach whose idea is borrowed from [Lom]. We need some terminology and conventions.

The algorithm transforms f step by step, and it may happen that some paths of the current multiflow are self-intersecting in edges (even if all paths in the initial multiflow are simple). By this reason, we should refine the definition of f^e in the Introduction as

$$(2.1) \quad f^e := \sum (\lambda_i n_i(e) : i = 1, \dots, k),$$

where $n_i(e)$ is the number of occurrences of e in P_i . We define $\Delta(e) = \Delta_f(e) := c(e) - f^e$ and call e *residual* if it is not saturated by f , i.e., if $\Delta(e) > 0$. Let $E^0 = E^0(f)$ be the set of residual edges in G . A *residual path* is a path in G whose all edges are residual. When it is not confusing, for a path $P = (x_0, e_1, x_1, \dots, e_k, x_k)$, we use the abbreviation $x_0 x_1 \dots x_k$. We say that P is an x_0 - x_k *path*, and denote a part of P from x_i to x_j by $P(x_i, x_j)$. Each path $P = x_0 \dots x_k$ in f is usually considered up to reversing (i.e., we do not distinguish between P and $P^{-1} = x_k x_{k-1} \dots x_0$). For $\{u, v\} \subseteq T$, the set of nodes occurred in the paths of the flow f_{uv} is denoted by V_{uv} (the *domain* of f_{uv}). Let $A = \{s_1, s_2\}$ and $B = \{t_1, \dots, t_r\}$ be the parts of H . The complete graph on T is denoted by $K_T = (T, E_T)$.

Each transformation of f never decreases the μ -value and any of the sets E^0 and V_{uv} , $uv \in E_T$; moreover, it increases at least one of these. Each iteration applies one of the seven operations below. The algorithm declares that the initial multiflow is not maximum and terminates when the μ -value increases (the *breakthrough* situation). This happens after performing the X-operation and, sometimes, the V-operation below. Also the breakthrough happens if there appears a residual path P connecting different terminals (for we can increase the μ -value by pushing some non-zero flow through P).

I-operation. Suppose there are a residual edge xy and a flow f_{uv} with V_{uv} containing x but not y . The *I-operation* chooses a path $P = x_0 \dots x_k$ in f_{uv} passing x , $x = x_i$ say, adds to f_{uv} the path $P' = x_0 \dots, x_i y x_i \dots x_k$ taken with a sufficiently small weight $\varepsilon > 0$ satisfying $\varepsilon < \lambda(P), \Delta(e)/2$, and accordingly reduces the weight of P by ε . This adds y to V_{uv} (while preserving the c -admissibility, the μ -value, and the sets E^0 and $V_{u'v'}$ for the other $u'v' \in E_T$).

V-operation. Suppose there are a u - v path $P = x_0 \dots x_k$ in f and a residual path $L = z_0 \dots z_d$ such that: z_0 is a terminal different from u and v ; $z_d = x_i$ for some i ; and either (a) $\mu(uz) + \mu(zv) > \mu(uv)$, or (b) for some $j \leq i$, the node x_j is not in V_{uz} , or both. The *V-operation* adds to f the u - z path $x_0 \dots x_i z_{d-1} \dots z_0$ and the z - v path $z_0 \dots z_d x_{i+1} \dots x_k$, each taken with a weight $\varepsilon > 0$ such that $\varepsilon < \lambda(P)$ and $\varepsilon < \Delta(e)/2$ for all edges e of L , and reduces $\lambda(P)$ by ε . This increases the μ -value, yielding the breakthrough, in case (a), and increases V_{uz} in case (b) (since x_j is added to V_{uz}).

O-operation. Suppose there are a u - v path $P = x_0 \dots x_k$ in f and a residual path $L = z_0 \dots z_d$ such that $z_0 = x_i$ and $z_d = x_j$ for some $0 \leq i < j \leq k$, and some edge e of the subpath $x_i \dots x_j$ is saturated. The *O-operation* adds to f the u - v path $x_0 \dots x_i z_1 \dots z_{d-1} x_j \dots x_k$ with a weight $\varepsilon > 0$ such that $\varepsilon < \lambda(P), \Delta(z_{p-1} z_p)$, $p = 1, \dots, d$, and reduces $\lambda(P)$ by ε . This increases E^0 (since e becomes residual).

Y-operation. Suppose there are a u - v path $P = x_0 \dots x_k$ and a u - v' path $Q = y_0 \dots y_q$ in f such that $v \neq v'$, $x_i = y_j$ for some $i, j > 0$, and some node of the subpath $y_0 \dots y_j$ is not in V_{uv} . The *Y-operation* adds to f the paths $P' = y_0 \dots y_j x_{i+1} \dots x_k$ and $Q' = x_0 \dots x_i y_{j+1} \dots y_q$, each taken with the same weight ε , $0 < \varepsilon < \lambda(P), \lambda(Q)$, and reduces each of $\lambda(P)$ and $\lambda(Q)$ by ε . This increases V_{uv} .

U-operation. Suppose there are a u - v path $P = x_0 \dots x_k$ and a u - v' path $Q = y_0 \dots y_q$ in f such that $x_i = y_j$ for some $i, j > 0$, $\mu(uv) + \mu(uv') = \mu(vv')$, and some edge e of the closed path $x_0 \dots x_i y_{j-1} \dots y_0$ is saturated. The *U-operation* adds to f the v - v' path $L = x_k x_{k-1} \dots x_i y_{j+1} \dots y_q$ with a weight ε , $0 < \varepsilon < \lambda(P), \lambda(Q)$, and reduce each of $\lambda(P)$ and $\lambda(Q)$ by ε . This increases E^0 (since e becomes residual), while preserving the μ -value.

Ψ -operation. Suppose there are a u - v path $P = x_0 \dots x_k$ and a z - w path $Q = y_0 \dots y_q$ in f such that: $x_i = y_j$ for some i, j ; the terminals v, z, w are different members of B ; u is in A ; and the subpath $x_0 \dots x_i$ contains a saturated edge e . Then $\mu(uv) = 1$ and $\mu(vz) = \mu(vw) = \mu(zw) = 2$. The *Ψ -operation* adds to f the v - z path $L = x_k \dots x_i y_{j-1} \dots y_0$ and the v - w path $L' = x_k \dots x_i y_{j+1} \dots y_q$, each with the same weight ε , $0 < \varepsilon < \lambda(Q), \frac{1}{2}\lambda(P)$, reduces $\lambda(P)$ by 2ε and reduces $\lambda(Q)$ by ε . This increases E^0 (since e becomes residual), and one can see that the μ -value preserves.

X-operation. Suppose there are a u - v path $P = x_0 \dots x_k$ and a z - w path $Q = y_0 \dots y_q$ in f such that: $x_i = y_j$ for some i, j ; u and z are different terminals in A ; and v and w are different terminals in B . The *X-operation* adds to f the u - z path $x_0 \dots x_i y_{j-1} \dots y_0$ and the v - w path $x_k \dots x_i y_{j+1} \dots y_q$, each with a weight ε , $0 < \varepsilon \leq \lambda(P), \lambda(Q)$, and reduces each of $\lambda(P), \lambda(Q)$ by ε . This increases the μ -value by 2ε (since $\mu(uv) = \mu(zw) = 1$ and $\mu(uz) = \mu(vw) = 2$), yielding the breakthrough.

The process terminates when the above operations are no longer applicable to the

current multiflow. Since each operation increases E^0 or V_{uv} for some $uv \in E_T$, the number of iterations is at most $|E| + |V||E_T|$. We show the following.

Lemma 2.1. *If neither the breakthrough happens nor any of the above operations is applicable, then the current multiflow f (as well as the initial one) is maximum.*

Proof. We first assume that $f_{s_1 t_p}$ is nonempty for each $p = 1, \dots, r$. Let S_1 be the node set of the component of (V, E^0) containing s_1 . For $p = 1, \dots, r$, define $V_p = V_{s_1 t_p}$ and $T_p = V_p - S_1$. Then $S_1 \cap T = \{s_1\}$ and $V_p \cap T = \{s_1, t_p\}$; for if S_1 contains a terminal $u \neq s_1$ or V_p contains a terminal $v \neq s_1, t_p$, we have the breakthrough (taking into account that $\mu(s_1 t_p) = 1 < \mu(s_1 v) + \mu(v t_p)$, whence the V-operation is applicable). Also $T_p \cap T_{p'} = \emptyset$ for $p \neq p'$ (otherwise we can apply the U-operation to an s_1-t_p path and an $s_1-t_{p'}$ path which share a common node in $T_p \cap T_{p'}$, increasing E^0). Therefore, the sets S_1, T_1, \dots, T_r and $S_2 = V - \cup(S_1 \cup T_1 \cup \dots \cup T_r)$ form a T -partition π of V ; let m be the $(2, r)$ -metric on V induced by π . We assert that

$$(2.2) \quad \text{for } e \in E, m(e) > 0 \text{ implies } \Delta(e) = 0,$$

and

$$(2.3) \quad \text{each } z-w \text{ path } Q = y_0 \dots y_q \text{ in } f \text{ is } m\text{-shortest, i.e., } m(Q) = \mu(zw).$$

Property (2.2) follows from the fact that all edges of the cut $\delta(S_1)$ are saturated, and similarly for the cuts $\delta(T_p)$ (for if e is a residual edge in some $\delta(T_p)$, then e belongs to $\delta(V_p)$, and we can apply the I-operation, increasing V_p). To see (2.3), consider possible cases (up to reversing Q). Note that $S_1 \subset V_p$ for each p (since the I-operation is impossible).

(a) Let $z = s_1$ and $w = t_p$. Then all nodes of Q are in V_p . Also Q meets $\delta(S_1)$ exactly once. Indeed, if $y_i \in S_1$ for some i , then the nodes y_0, y_1, \dots, y_i are contained in S_1 (otherwise E^0 can be increased by use of the O-operation). Hence, $m(Q) = 1 = \mu(s_1 t_p)$.

(b) Let both z, w be in B , $z = t_1$ and $w = t_2$ say. Let y_i be the last node in T_1 , and y_j the first node in T_2 . Then $y_0, \dots, y_i \in T_1$ and $y_j, \dots, y_q \in T_2$. Indeed, suppose $y_{j'}$ is not in T_1 for some $i' < i$. If $y_{i'}$ is in S_1 , then the V-operation applied to Q and a residual path from s_1 to $y_{i'}$ increases V_2 by y_i (as $y_i \in T_1$ implies $y_i \notin V_2$). And if $y_{i'} \notin S_1$, then $y_{i'} \notin V_1$ and, therefore, the Y-operation applied to Q and a t_1-s_1 path in f that passes y_i increases V_1 by $y_{i'}$. Now consider the nodes y_{i+1}, \dots, y_{j-1} . None of them is in T_p for $p \neq 1, 2$ (otherwise the Ψ -operation for Q and an s_1-t_p path in f containing such a node increases V_1). If these nodes are entirely contained in one of S_1 and S_2 , then, assuming $j > i + 1$, we have $m(Q) = m(y_i y_{i+1}) + m(y_{j-1} y_j) = 1 + 1 = 2 = \mu(t_1 t_2)$, as required (in case $j = i + 1$ we also have $m(Q) = 2$). Now suppose that $y_g \in S_1$ and

$y_h \in S_2$ for some $i < g < h < j$ (case $i < h < g < j$ is symmetric). Let P be a t_2 - s_1 path in f which contains y_g . Then the Y-operation for P and Q increases V_2 by y_h .

(c) Let $z = s_2$ and $w = t_p$. Then Q does not meet $V_{p'}$ for $p' \neq p$ (otherwise we can apply the X-operation, increasing the μ -value). So all nodes of Q are in $T_p \cup S_2$. Also $y_i \in S_2$ implies $y_0, \dots, y_i \in S_2$ (otherwise one can apply the Y-operation, increasing V_p). Hence, Q passes exactly one edge between S_2 and T_p , yielding $m(Q) = 1 = \mu(zw)$.

(d) Let $z = s_1$ and $w = s_2$. Take the last node y_i of Q not in S_2 . Then $y_i \in V_p$ for some p , whence $y_0, \dots, y_i \in V_p$ (otherwise one can increase V_p by use of the Y-operation). Now take the last node of Q in S_1 . Then $y_0, \dots, y_j \in S_1$ (otherwise one can increase E^0 by use of the O-operation). Now $i = j$ gives $m(Q) = m(y_i y_{i+1}) = 2$, and $i > j$ gives $m(Q) = m(y_j y_{j+1}) + m(y_i y_{i+1}) = 1 + 1 = 2$. Thus, $m(Q) = \mu(zw)$.

In a general case, for each p with $f_{s_1 t_p}$ empty, we define T_p to be the node set of the component of (V, E^0) containing t_p . Again one can see that $\{S_1, S_2, T_1, \dots, T_r\}$ is a T -partition of V . We leave it to the reader, as an exercise, to check that (2.2)–(2.3) hold in this case too (by arguing as above or, sometimes, even simpler).

Now (2.2) and (2.3) (being, in fact, the complementary slackness conditions for (1.4) and its dual problem) imply $\langle \mu, f \rangle = c \cdot m$. Indeed, let f consist of the u_i - v_i paths Q_i , $i = 1, \dots, q$, whose weights $\lambda_1, \dots, \lambda_q$ are nonzero. Then

$$\begin{aligned} \langle \mu, f \rangle &= \sum_{i=1}^q \lambda_i \mu(u_i v_i) \leq \sum_{i=1}^q \lambda_i m(Q_i) \\ &= \sum_{e \in E} f^e m(e) \leq \sum_{e \in E} c(e) m(e) = c \cdot m, \end{aligned}$$

and equality holds throughout in this expression because of (2.2) and (2.3).

Thus, f is a maximum multiflow and m is a minimum $(2, r)$ -metric. •

Let k be the number of paths of the initial multiflow f . W.l.o.g. one may assume that the length (number of edges) of each of these paths is $O(|V|)$. Since each operation applied creates at most two new paths, the number of paths of each current multiflow in the above process is polynomial in $|V|, |E|, k$. Moreover, after each iteration we can rearrange paths of the current multiflow f so as to make the length of each path in f be $O(|V|)$, without decreasing the μ -value and the sets E^0 and V_{uv} , $uv \in E_T$. Then each of the above operations can be performed in time polynomial in $|V|, |E|, k$, whence the running time of the maximality check-up algorithm is polynomial in $|V|, |E|, k$ as well.

Remark. In fact, in the above process it suffices to apply only those operations that increase the μ -value or the set E^0 or some set $V_{s_i t_j}$ (rather than $V_{s_1 s_2}$ or $V_{t_i t_j}$). One can check that Lemma 2.1 remains valid in this case too. Such a modification is more

efficient because it applies only $O(|E| + |T||V|)$ operations instead of $O(|E| + |T|^2|V|)$ operations in the algorithm described above.

The analysis of the above algorithm shows that the weights of paths and the values of residual capacities are, in essence, not important. The algorithm can maintain only the set of paths of the current multiflow and the set E^0 of residual edges, and perform each iteration in terms of these two sets. In particular, this implies the following important fact (it will be used later).

Corollary 2.2. *Let $f = (P_1, \dots, P_k; \lambda_1, \dots, \lambda_k)$ be a multiflow for G, c, T , and let E^0 be the set of residual edges for c and f . Let f' be the multiflow consisting of the same paths P_1, \dots, P_k but taken with weight one each, and let $c' = \chi^{P_1} + \dots + \chi^{P_k} + \chi^{E^0}$. Then f is maximum for G, c, μ if and only if f' is maximum for G, c', μ . •*

3. Integer augmentation

In this section we deal with a special case when $G = (V, E)$ is an *inner Eulerian graph* (i.e., all inner nodes $x \in V - T$ have even degrees) and all capacities $c(e)$ of edges $e \in E$ are ones. We allow G to have multiple edges but not loops and denote $\nu(G, c, \mu)$ by $\nu(G, \mu)$. As before, $T = \{s_1, s_2, t_1, \dots, t_r\}$ and $\mu = d^H$, where $H = (T, U)$ is the graph $K_{2,r}$ with the parts $A = \{s_1, s_2\}$ and $B = \{t_1, \dots, t_r\}$. In the input of the integer augmentation problem, we are given an *integer* multiflow f in (G, T, c) whose μ -value is not maximum, and the goal is to augment its μ -value by at least one, i.e., to find an integer multiflow f' with $\langle \mu, f' \rangle > \langle \mu, f \rangle$. We describe an algorithm for finding such an f' , in time polynomial in $|V|, |E|$.

Since all capacities are ones, one may assume that every integer multiflow is a set of pairwise edge-disjoint T -paths (which have unit weights). Let E^0 be the sets of residual (not used in the paths of f) edges of G . One may assume that each component of (V, E^0) contains at most one terminal; otherwise the problem is trivial. Then E^0 is representable as the union of pairwise edge disjoint circuits. In what follows by a *decomposition* we mean a decomposition \mathcal{D}' of G (or another inner Eulerian graph in question) into T -paths and circuits. The μ -value $\langle \mu, \mathcal{D}' \rangle$ of \mathcal{D}' is the μ -value of the corresponding multiflow formed by the T -paths in \mathcal{D}' .

Let \mathcal{D} be a decomposition including the initial f , and let $\bar{\nu} = \langle \mu, f \rangle = \langle \mu, \mathcal{D} \rangle$. The algorithm we develop handles \mathcal{D} rather than f , attempting to transform \mathcal{D} into a decomposition \mathcal{D}' with a greater μ -value.

To simplify the algorithm description, we assume that each inner node of G has

degree at most four. This leads to no loss of generality. Indeed, denote by $E(x)$ the set of edges incident to a node x , and suppose that $|E(x)| > 4$ for some $x \in V - T$. Then we can transform the graph at x as follows. Partition each edge $e = xy$ into two edges xz_e and $z_e y$ in series, and connect the nodes $z_e, e \in E(x)$, by a simple circuit C of new (residual) edges. For each pair $e, e' \in E(x)$ such that e, x, e' are consecutive elements of some member of \mathcal{D} , replace $xz_e, xz_{e'}$ by a single edge $z_e z_{e'}$, and then remove x . See Fig. 2 for an illustration. This transformation creates $|E(x)|$ nodes with degree four in place of x , and \mathcal{D} is transformed, in a natural way, into a decomposition \mathcal{D}' with the same μ -value in the resulting graph G' . Note that \mathcal{D}' is not maximum in G' . (For otherwise take an optimal T -partition π' for G' . Then the nodes of C are entirely contained in one member of π' . This implies that π' induces an optimal T -partition π for G , whence \mathcal{D} is maximum; a contradiction.) Note also that any integer multiflow g' in G' can be easily transformed into an integer multiflow f' with the same μ -value in G . So if we succeed to find g' in G' with $\langle \mu, g' \rangle > \langle \mu, g \rangle$, it gives a multiflow f' in G with $\langle \mu, f' \rangle > \bar{v}$.




Fig. 2

Let W be the set of inner nodes of degree four in G . For each $x \in W$, \mathcal{D} induces in a natural way a *bi-partition* $\rho(x) = \rho^{\mathcal{D}}(x)$ of the set $E(x) = \{e_1, e_2, e_3, e_4\}$ into two pairs, $\{e_1, e_2\}$ and $\{e_3, e_4\}$ say; we write $\rho(x) = \{e_1, e_2 | e_3, e_4\}$. W.l.o.g., one may assume that each path or circuit in \mathcal{D} is simple and that no path P in \mathcal{D} contains a terminal as an intermediate node (otherwise split P at such a node into two T -paths; this does not decrease the μ -value). We say that such a \mathcal{D} is *simple*. The non-maximality of f and the fact that no residual path connects different terminals imply that W is nonempty. Note that \mathcal{D} is determined uniquely by the bi-partitions $\rho(x)$ for $x \in W$. Moreover, every set of bi-partitions $\rho'(x)$ at the elements $x \in W$ determines uniquely a (possibly non-simple) decomposition $\mathcal{D}_{\rho'}$ in which each circuit contains at most one terminal and each T -path meets T only at its ends.




Fig. 3

Consider $x \in W$, and let $\rho(x) = \{e_1, e_2 | e_3, e_4\}$. The *splitting operation* at x with respect to a bi-partition $\bar{\rho} = \{e_i, e_j | e_p, e_q\}$ of $E(x)$ (possibly $\bar{\rho} = \rho(x)$) transforms G into a graph G' by replacing the end x of e_i, e_j by a new node x' and replacing the end x of e_p, e_q by a new node x'' ($x' \neq x''$); see Fig. 3. This gives a (possibly non-simple) decomposition $\mathcal{D}' = \mathcal{D}_{\rho'}$, where $\rho'(x) = \bar{\rho}$ and $\rho'(y) = \rho(y)$ for all $y \in W - \{x\}$. We call such an operation

- (i) an *augmenting* splitting if $\langle \mu, \mathcal{D}' \rangle > \bar{\nu}$;
- (ii) a *feasible* splitting if $\langle \mu, \mathcal{D}' \rangle = \bar{\nu}$;
- (iii) a *good* splitting if it is feasible and \mathcal{D}' is still not maximum in the new graph G' , i.e., $\nu(G', \mu) > \bar{\nu}$;
- (iv) a *laminar* splitting if $\bar{\rho} = \rho(x)$, and *cross-splitting* otherwise.

In particular, the laminar splitting is, obviously, feasible (but may not be good). Clearly to compute the value $\langle \mu, \mathcal{D}' \rangle$ and compare it with $\bar{\nu}$ is easy. If there is an augmenting splitting at some node $x \in W$, we immediately obtain an integer multifold with a greater μ -value; so assume this is not the case. Also using the maximality check-up algorithm from the previous section, we can examine all feasible splittings at the nodes in W to seek a good one among them, and once a good splitting is found, we transform G into the graph G' whose set W' of inner nodes of degree four is smaller than W .

Assume that the laminar splitting is not good at each node in W . We will rely on the following simple fact.

Statement 3.1. For $x \in W$, if both cross-splittings at x are feasible, then at least one of them is good.

Proof. Take a decomposition \mathcal{D}' of G with $\langle \mu, \mathcal{D}' \rangle > \bar{\nu}$ (it exists by Theorem 1.1 since f is not maximum in G), and let G' be obtained by the splitting at x w.r.t. the bi-partition $\rho' = \rho^{\mathcal{D}'}(x)$. Then $\nu(G', \mu) \geq \langle \mu, \mathcal{D}' \rangle > \bar{\nu}$. Since the laminar splitting at x is not good, $\rho' \neq \rho^{\mathcal{D}'}(x)$. Hence, with respect to \mathcal{D} , the given splitting is a good cross-splitting (as it is feasible). •

In what follows we assume that no feasible splitting at any node is good. Let P and P' be the members of \mathcal{D} passing $x \in W$. The above statement enables us to eliminate the following situations in which each of the two cross-splittings at x is feasible or even augmenting:

- (3.1) (i) at least one of P, P' is a circuit;
- (ii) P is a u - v path and P' is a z - w path and each of $\mu(uz) + \mu(vw)$ and $\mu(uw) +$

$\mu(vz)$ is greater than or equal to $\mu(uv) + \mu(zw)$.

As before, we consider each path in f up to reversing, and for $u, v \in T$, denote by V_{uv} the domain of the flow f_{uv} . The impossibility of (3.1) implies:

- (3.2) all members of \mathcal{D} are (simple) T -paths;
- (3.3) for $i = 1, 2$ and $j = 1, \dots, r$, if V_{s_i, t_j} and V_{uv} with $\{s_i, t_j\} \neq \{u, v\}$ share an inner node, then $\{u, v\}$ is either $\{s_i, s_{i'}\}$ or $\{t_j, t_{j'}\}$; in particular, the flows f_{s_i, t_j} are pairwise openly disjoint, i.e., $V_{s_i, t_j} \cap V_{s_{i'}, t_{j'}} = \{s_i, t_j\} \cap \{s_{i'}, t_{j'}\}$ for $(i, j) \neq (i', j')$.

Note that f contains at least one s_1-s_2 or t_i-t_j path (otherwise (3.3) would imply that $W = \emptyset$, whence f is already maximum). Until now we have tried to get the desired augmentation or decrease the set W by applying a single splitting operation, and (3.2)–(3.3) expose all we are able to get on this way. However, one can rearrange the decomposition \mathcal{D} more globally by combining splittings at several nodes at once (without decreasing the μ -value) in order to get a situation as in (3.1) and then make a crucial splitting.

For example, let f contains an s_1-t_1 path $P = x_0 \dots x_k$, an s_1-t_2 path $P' = y_0 \dots y_p$ and an s_1-s_2 path $P'' = z_0 \dots z_q$ such that P'' meets both P and P' at intermediate nodes, $x_i = z_j =: x$ and $y_{i'} = z_{j'} =: y$ for $j < j'$ say. Then no single splitting at x or at y is good. Nevertheless, $P \cup P' \cup P''$ is decomposed into the s_1-s_2 path $Q = y_0 \dots y_{i'} z_{j'+1} \dots z_q$, the t_1-t_2 path $Q' = x_k \dots x_i z_{j+1} \dots z_{j'} y_{i'+1} \dots y_p$ and the circuit $C = x_0 \dots x_i z_{j-1} \dots z_0$. Hence, replacing P, P', P'' in \mathcal{D} by Q, Q', C makes \mathcal{D}' with $\langle \mu, \mathcal{D}' \rangle = \langle \mu, \mathcal{D} \rangle$, and now a good splitting at x becomes possible since x belongs to a T -path and a circuit.

In general, we rearrange f as follows. Note that at least one of the flows f_{s_i, t_j} , f_{s_1, t_1} say, is nonzero (otherwise f is, obviously, maximum). We construct a minimal set $L = L(s_1, t_1)$ such that $V_{s_1, t_1} \subseteq L$ and each s_1-s_2 or t_1-t_p path meets the cut $\delta(L)$ precisely once. Such an L is unique and is constructed by use of a labelling method. Initially, set $L := V_{s_1, t_1}$. Then L is increased step by step by the following rule:

- (3.4) choose an s_1-s_2 or t_1-t_p path $P = x_0 \dots x_k$ in f such that for some $0 < i < j < k$, $x_i \notin L$ and $x_j \in L$, and update $L := L \cup \{x_0, \dots, x_j\}$.

The process terminates when L cannot be increased by (3.4). Clearly $L \cap T = \{s_1, t_1\}$. Let $U_1 = \{\{s_1, s_2\}, \{t_1, t_2\}, \{t_1, t_3\}, \dots, \{t_1, t_r\}\}$ and $\bar{U}_1 = U - (U_1 \cup \{s_1, t_1\})$. Suppose that there is a node $x \in L - \{s_1, t_1\}$ such that

- (3.5) x belongs to a set V_{zw} , where $\{z, w\} \in \bar{U}_1$.

We show that the submultiflow g in f that consists of the flows $f_{s_1 t_1}$ and f_{uv} , $\{u, v\} \in U_1$ can be rearranged within L so that there appears an s_1 - t_1 path P containing x . Then P and the path in f_{zw} containing x , where $\{z, w\}$ is as in (3.5), are in situation (3.1)(ii) and, therefore, one can apply an augmenting or good splitting at x . The task of finding such a rearrangement is reduced to the multiflow demand problem (defined in the Introduction) for the graph $\Gamma = (F, Z)$, demand pairs D and demands d on the values of flows connecting pairs in D , where:

- (i) Γ is formed by the nodes and edges occurring in g or in T ;
- (ii) D consists of the pairs $\{s_1, x\}$, $\{x, t_1\}$, $\{s_1, t_1\}$ and the pairs in U_1 ;
- (iii) $d(s_1 x) = d(x t_1) = 1$, $d(s_1 s_2) = |f_{s_1 s_2}| - 1$ and $d(uv) = |f_{uv}|$ for $\{u, v\} \in U_1$ ($|f_{uv}|$ denotes the number of paths in f_{uv}).

Statement 3.2. *The above demand problem has an integer solution, i.e., there are pairwise edge-disjoint T' -paths Q_1, \dots, Q_q in Γ such that each pair $\{u, v\}$ in D is connected by $d(uv)$ of these paths, where $T' = T \cup \{x\}$.*

Proof. For $X \subseteq F$, define d_X as in (1.7), and let $\Delta(X) = |\delta^\Gamma(X)| - d_X$. It is easy to see that $\Delta(X)$ is an even integer for any X . Also each pair in D meets at least one of s_1, t_1 , i.e., the graph induced by D is the union of two stars. Hence, the problem has an integer solution provided that the cut condition holds, i.e., if $\Delta(X) \geq 0$ for all $X \subseteq F$ (by a reduction of Dinitz (see [ADK]) from the two-star commodity demand problem to the two commodity one and by a theorem of Rothschild and Whinston concerning the latter [RW]). Suppose that $\Delta(X) < 0$ for some X ; let for definiteness $s_1 \in X$. Considering the feasible multiflow g in Γ , we observe that $\Delta(X) < 0$ is possible only if $s_1, t_1 \in X$ and $x \notin X$. One may assume that $X \cap T' = \{s_1, t_1\}$ (for if $v \in X$ for some $v \in T - \{s_1, t_1\}$, then, obviously, $\Delta(X - \{v\}) \leq \Delta(X)$).

Let $\bar{d} = \sum(|f_{uv}| : \{u, v\} \in U_1)$. Then $|\delta^\Gamma(X)| \geq \bar{d}$ and $\bar{d} = d_X - 2$. Therefore, the evenness of $\Delta(X)$ implies $\Delta(X) = -2$, whence $|\delta^\Gamma(X)| = \bar{d}$. The latter equality implies that each path in each f_{uv} , $\{u, v\} \in U_1$, meets $\delta^\Gamma(X)$ exactly once. Then no node outside X can be labelled by rule (3.5). So $x \notin L$; a contradiction. •

Let Q_i 's be as in Statement 3.2, and let for definiteness Q_1 and Q_2 be the paths from s_1 to x and from x to t_1 , respectively. Then the multiflow g' formed by the paths Q_3, \dots, Q_q and the s_1 - t_1 path being the concatenation of Q_1 and Q_2 satisfies $\langle \mu, g' \rangle = \langle \mu, g \rangle$. We replace in f the part g by g' , obtaining situation (3.1)(ii), and proceed with an augmenting or good splitting at x .

If there is no x as in (3.5), we consecutively construct the labelled sets $L_2 = L(s_1, t_2), \dots, L_r = L(s_1, t_r)$ in a similar way. At least one set $L_i - \{s_1, t_i\}$ must meet

V_{uv} with $\{u, v\}$ different from $\{s_1, t_i\}, \{s_1, s_2\}, \{t_i, t_q\}, q \in \{1, \dots, r\} - \{i\}$ (otherwise, in view of (3.3), $L_i \cap L_j = \{s_1\}$ for all $i \neq j$, whence the sets $\{s_1\}, L_i - \{s_1\}$ ($i = 1, \dots, r$) and $V - (L_1 \cup \dots \cup L_r)$ form an optimal T -partition, implying that f is maximum).

The above demand problem can be solved in time polynomial in $|V|, |E|$ (e.g., by the method behind the proof in [RW]). Since the number of splittings we apply does not exceed $|V|$, the whole time needed to find an integer augmentation of f is polynomial.

4. Scaling method

In this section we put together the above arguments to design a polynomial algorithm for solving (1.6) and the corresponding integer multiflow problem (1.3). As before, μ is the path metric of the graph $K_{2,r}$ and the capacity function c on the edges of $G = (V, E)$ is assumed to be inner Eulerian.

At the high level, the algorithm applies a capacity scaling approach and consists of *big* (or scaling) iterations. The number of these iterations is equal to the size of the largest capacity $\|c\|$ in binary notation.

More precisely, let I be $\lceil \log_2(\|c\| + 1) \rceil$. For $i = 0, \dots, I$ and $e \in E$, define the truncated capacity $c_i(e) = \lfloor c(e)/2^{I-i} \rfloor$. Then $c_0 = 0$ and $c_I = c$. In the input of i th big iteration, there is a maximum *half-integer* multiflow g_{i-1} for G, c_{i-1}, μ (letting g_0 be zero multiflow), and the goal is to find a maximum half-integer multiflow g_i for c_i . (The reason why we are forced to deal with half-integer multiflows is that c_i needs not be inner-Eulerian.) The final, I th, big iteration will find a maximum *integer* multiflow for G, c, μ along with a minimum $(2, r)$ -metric.

We describe i th iteration, $i < I$. It considers $c' = 2c_i$ as the capacity function and $g' = 4g_{i-1}$ as the initial multiflow (i.e., the weights of all paths in g_{i-1} are increased by a factor of four). Then c' is inner Eulerian and g' is c' -admissible and integral. Moreover, we observe that

$$(4.1) \quad \tau(c') - \langle \mu, g' \rangle \leq 4|E|,$$

where $\tau(c')$ stands for $\tau(G, c', \mu)$. Indeed, let m be a minimum $(2, r)$ -metric for c_{i-1} . Then g' and m are optimal for $4c_{i-1}$, i.e., $\langle \mu, g' \rangle = (4c_{i-1}) \cdot m$. For each $e \in E$, we have $m(e) \leq 2$ and $c'(e) = 2c_i(e) \leq 4c_{i-1}(e) + 2$. Therefore, $\tau(c') \leq c' \cdot m \leq (4c_{i-1}) \cdot m + 4|E|$, yielding (4.1).

Thus, g' is nearly optimal for c' , and it takes at most $4|E|$ integer augmentations to transform g' into a maximum integer multiflow for c' . However, the integer augmentation algorithm from Section 3 is sensitive to the capacity values (it is polynomial only for small capacities and pseudo-polynomial in general). To this reason, every time

we turn to this algorithm, we first transform c' and the current integer multiflow g as follows.

Let $g = (P_1, \dots, P_k; \lambda_1, \dots, \lambda_k)$, where all λ_i 's are positive integers. One may assume that each flow g_{uv} in g consists of at most $|E|$ paths (for we can rearrange g_{uv} , if needed, using standard flow decomposition techniques [FF]). Then k is $O(|T|^2|E|)$. We replace g by the multiflow h consisting of the same paths P_1, \dots, P_k but taken with weight one each. Also we represent the residual capacities $\Delta(e) = c'(e) - g^e$ as $\Delta = \alpha_1 \chi^{C_1} + \dots + \alpha_q \chi^{C_q}$, where $q \leq |E|$, each α_i is a positive integer and each C_i is a circuit in G (χ^C is the incidence vector of a circuit C in \mathbb{R}^E). We replace c' by $c'' = \chi^{P_1} + \dots + \chi^{P_k} + \chi^{C_1} + \dots + \chi^{C_q}$. Then $\|c''\|$ is only $O(|T|^2|E|)$ and, therefore, replacing each edge e by $c''(e)$ parallel edges makes an inner Eulerian multigraph whose size is polynomial in $|V|, |E|$. By Corollary 2.2, h is not maximum for c'' if and only if g is not maximum for c' . Moreover, an integer augmentation of h by use of the algorithm in Section 3 determines an integer augmentation of g in a natural way.

Summing up the above arguments, we conclude that each big iteration is performed in strongly polynomial time and finds a maximum integer multiflow g for $2c_i$, giving the maximum half-integer multiflow $g_i = \frac{1}{2}g$ for c_i . The final, I th, iteration is applied to the inner Eulerian $c_I = c$ and integer multiflow $g' = 2g_{I-1}$ (instead of $2c_I$ and $4g_{I-1}$) and finds a maximum integer multiflow f for the initial G, c, μ . Also the last application of the algorithm from Section 2 constructs a minimum $(2, r)$ -metric m . Thus, the total time of our algorithm is polynomial in $|V|, |E|$ and linear in $\log \|c\|$, as required.

5. Algorithm for sparse frames

We will use some properties of modular graphs. A graph $H = (T, U)$ is called *modular* if each three nodes $v_0, v_1, v_2 \in T$ have a *median*, i.e., a node $z \in T$ such that $d^H(v_i, z) + d^H(z, v_j) = d^H(v_i, v_j)$ for all $0 \leq i < j \leq 2$. If every isometric subgraph of H is modular, then H is said to be *hereditary modular*. It is easy to see that any modular graph is bipartite.

Bandelt [B88] proved the following theorem: a bipartite graph H has no isometric k -circuit with $k \geq 6$ if and only if H is hereditary modular. Thus, the frames (figured in Theorem 1.2) are precisely the orientable hereditary modular graphs. Modular graphs have the following property (see [B85]):

- (5.1) for any orbit Q of a modular graph H and any $u, v \in T$, if P is a shortest u - v path and P' is a u - v path, then $|P \cap Q| \leq |P' \cap Q|$.

For a function $\ell : U \rightarrow \mathbb{R}_+$ on the edges of a graph $H = (T, U)$, we denote by $d^{H, \ell}$

the path metric for (H, ℓ) , i.e., $d^{H, \ell}(xy)$ is the minimum ℓ -length $\ell(P) = \sum(\ell(e_i) : i = 1, \dots, k)$ of a path $P = (x_0, e_1, x_1, \dots, e_k, x_k)$ connecting nodes x and y in H . An ℓ -shortest path is a path shortest for the metric $d^{H, \ell}$. We say that ℓ is *orbit-invariant* if it is constant within each orbit of H . Consider a modular graph $H = (T, U)$, and let Q_1, \dots, Q_k be the orbits of H . From (5.1) it follows that

(5.2) for any nonnegative orbit-invariant function ℓ on U , each shortest path in H is ℓ -shortest.

Indeed, let $h_i = \ell(e)$ for $e \in Q_i$. For two u - v paths P and P' in H , we have $|P| = n_1 + \dots + n_k$, $|P'| = n'_1 + \dots + n'_k$, $\ell(P) = h_1 n_1 + \dots + h_k n_k$ and $\ell(P') = h_1 n'_1 + \dots + h_k n'_k$, where $|L|$ is the number of edges of a path L , $n_i = |P \cap Q_i|$ and $n'_i = |P' \cap Q_i|$. If P is shortest, then $n_i \leq n'_i$ for each i (by (5.1)) implies $\ell(P) \leq \ell(P')$ since ℓ is nonnegative.

For $i = 1, \dots, k$, define ℓ_i to be the incidence vector of Q_i (i.e., $\ell_i(e) = 1$ if $e \in Q_i$ and 0 if $e \in U - Q_i$), and let $\mu_i = d^{H, \ell_i}$.

Statement 5.1. $d^H = \mu_1 + \dots + \mu_k$.

Proof. Consider a shortest u - v path P in H . By (5.2), P is ℓ_i -shortest for $i = 1, \dots, k$. Therefore, $|P| = \ell_1(P) + \dots + \ell_k(P)$ implies $d^H(uv) = \mu_1(uv) + \dots + \mu_k(uv)$. •

In what follows we assume that the above graph H is a sparse frame. For $i = 1, \dots, k$, let $H_i = (T_i, U_i)$ be the graph in the definition of sparse frames in the Introduction. Define π_i to be the partition of T formed by the node sets of components of the graph $(T, U - Q_i)$. We identify each node of H_i with some node in the corresponding member of π_i . Then $T_i \subseteq T$ and μ_i is a 0-extension of $\hat{\mu}_i = d^{H_i}$ to T .

Consider a graph $G = (V, E)$ with $V \supseteq T$ and a capacity function c on E . Let $G_i = (V_i, E)$ be the graph (with possible parallel edges and loops) obtained from G by shrinking each set in π_i into the corresponding node of H_i . Then every 0-extension m' of μ_i to V one-to-one corresponds, in a natural way, to a 0-extension of $\hat{\mu}_i$ to V_i , denoted by $\gamma_i(m')$; obviously, $c \cdot m' = c \cdot (\gamma_i(m'))$. Let $\tau_i = \tau(G_i, c, \hat{\mu}_i)$.

Let m be a minimum 0-extension for G, c, μ , i.e., $c \cdot m = \tau$ ($= \tau(G, c, \mu)$). From Statement 5.1 it follows that m is represented as $m = m_1 + \dots + m_k$, where each m_i is a 0-extension of μ_i to V . This implies

$$(5.3) \quad \tau \geq \tau_1 + \dots + \tau_k.$$

Moreover, (5.3) turns into equality for any frame H . Indeed, for $i = 1, \dots, k$, take a 0-extension \hat{m}_i of $\hat{\mu}_i$ to V_i with $c \cdot \hat{m}_i = \tau_i$, and let m_i be the corresponding 0-extension $\gamma^{-1}(\hat{m}_i)$ of μ_i to V . Then $m = m_1 + \dots + m_k$ is an extension of μ to V , whence

$c \cdot m \geq \tau^*$ ($= \tau^*(G, c, \mu)$). This implies $\tau^* \leq \tau_1 + \dots + \tau_k$, yielding $\tau \leq \tau_1 + \dots + \tau_k$, in view of $\tau = \tau^*$.

The above arguments prompt an approach to solve (1.1) in our case. First we find the numbers τ_i by use of the above algorithm for the minimum $(2, r)$ -metric problem; their sum is just τ . (Note that if H_i is $K_{1,r}$, the problem for $G_i, c, \hat{\mu}_i$ is reduced to r minimum cut computations. In fact, one can show that $r = 1$ is only possible but this is not important for us.) In order to find the desired minimum 0-extension of μ to V , we can apply a method suitable for arbitrary frames (see [K98]). More precisely, choose a node $x \in V - T$ and terminal $t \in T$ and compute $\tau' = \tau(G', c, \mu)$, where G' is obtained from G by identifying x and t . If $\tau' = \tau$, we replace G by G' . And if $\tau' > \tau$, choose another terminal t' to be identified with x and do similarly, and so on (since μ is minimizable, at least one choice of a terminal t for x gives $\tau' = \tau$). Then repeat the procedure with a next node $x' \in V - T$. Eventually, after at most $|V - T||T|$ iterations, we obtain a graph $\tilde{G} = (\tilde{V}, E)$ with $\tilde{V} = T$ and $\tau(\tilde{G}, c, \mu) = \tau$, which determines the minimum 0-extension for G, c, μ in an obvious way.

However, we can suggest a more efficient method in which the problem with the same μ_i is solved only once. We say that a metric m on V is *cyclically even* if $m(xy) + m(yz) + m(zx)$ is an even integer for any $x, y, z \in V$ (in particular, m is integral). An extension m of μ to V is called *tight* if each two $x, y \in V$ belong to an m -shortest T -path, i.e., $m(sx) + m(xy) + m(yt) = \mu(st)$ for some $s, t \in T$.

As before, we first find an optimal 0-extension m_i of μ_i to V for each i and put $m = m_1 + \dots + m_k$. Each m_i is cyclically even (as H_i is bipartite), therefore, m is cyclically even as well. Next we find a cyclically even tight extension $m' \leq m$ of μ (note that m may not be tight). Initially, set $m' = m$. Choose $x, y \in V$ such that $m'(xy) \geq 2$ and

$$\Delta(xy) := \min\{m'(sx) + m'(xy) + m'(yt) - \mu(st) : s, t \in T\} > 0:$$

Put $\ell(xy) = \max\{0, m'(xy) - \Delta(xy)\}$ and $\ell(e) = m'(e)$ for each $e \in E_V - \{xy\}$, and update $m' := d^{K, \ell}$, where $K = (V, E_V)$ is the complete graph on V . Then any T -path P in K satisfies $\ell(P) \geq \mu(st)$, where s, t are the ends of P , i.e., the new m' is an extension of μ . Also one can see that m' is cyclically even. We do similarly for a next pair $x', y' \in V$, and so on. After $O(|V|^2)$ iterations we obtain a cyclically even extension m' of μ such that for any $x, y \in V$,

$$(5.4) \quad \text{if } m'(xy) \geq 2, \text{ then } \Delta(xy) = 0.$$

Statement 5.2. $\Delta(xy) = 0$ for all $x, y \in V$.

Proof. By (5.4), it suffices to examine a pair x, y with $m'(xy) \leq 1$. Let $m'(xy) = 1$. Choose $z \in V$ such that both $m'(zx), m'(zy)$ are nonzero (if no such z exists, then

$m'(sx) = m'(ty) = 0$ for some $s, t \in T$, whence $\Delta(xy) = 0$). Since m' is cyclically even and $m'(xy) = 1$, either $m'(zx) + m'(xy) = m'(zy)$ or $m'(zy) + m'(yx) = m'(zx)$. Assume the former (the latter is similar). Then $m'(zy) \geq 2$. Therefore, there exist $s, t \in T$ such that $m'(sz) + m'(zy) + m'(yt) = \mu(st)$. This implies $m'(sx) + m'(xy) + m'(yt) = \mu(st)$, yielding $\Delta(xy) = 0$. In case $m'(xy) = 0$, $\Delta(xy) = 0$ follows from $\Delta(xz) = 0$ for $z \in V$ with $m'(xz) > 0$. •

Thus, m' is tight. Also $m' \leq m$ implies $c \cdot m' \leq c \cdot m$, whence $c \cdot m' = \tau$. We assert that m' is a 0-extension of μ and, therefore, m' is an optimal solution to (1.1) in our case. The proof relies on a result concerning the tight spans for frames. More precisely, it is shown in [Dr,Is] that for every metric space (X, d) , there exists a unique tight extension (\mathcal{X}, δ) such that any tight extension (X', d') of (X, d) is isometrically embeddable in (\mathcal{X}, δ) , i.e., there is a mapping $\gamma : X' \rightarrow \mathcal{X}$ satisfying $\gamma(s) = s$ for all $s \in X$ and $d'(xy) = \delta(\gamma(x)\gamma(y))$ for all $x, y \in X'$. Such an (\mathcal{X}, δ) is called the *tight span* (or injective envelope, or T_X -space) of (X, d) . [K98] gives an explicit combinatorial construction for the tight span of the path metric of a frame. The proof of validity of this construction includes the following result.

Statement 5.3 [K98]. *Let $H = (T, U)$ be a frame and let m' be a tight extension of d^H to V . For each $x \in V$, at least one of the following is true:*

- (i) $m'(vx) = 0$ for some $v \in T$;
- (ii) $m'(ux) + m'(xv) = 1$ for some edge $uv \in U$;
- (iii) $m'(v_0x) + m'(xv_2) = m'(v_1x) + m'(xv_3) = 2$ for some 4-circuit $C = v_0v_1v_2v_3v_0$ of H .

Using this result, we show that for each $x \in V$, there exists $t \in T$ with $m'(tx) = 0$. This is immediate in cases (i) and (ii) of Statement 5.3. So assume that we are in case (iii) and $m'(v_i x) > 0$ for $i = 0, 1, 2, 3$. Then $m'(v_i x) = 1$ for each i . But $m'(v_0x) + m'(xv_1) + m'(v_1v_0) = 1 + 1 + 1 = 3$, contrary to the fact that m' is cyclically even. Thus, m' is indeed a 0-extension.

This gives a “purely combinatorial” algorithm for solving (1.1) with μ to be the path metric of a sparse frame H . This algorithm involves $O(|T|)$ minimum $(2, r)$ -metric computations and runs in time polynomial in $|V|, |E|$ and linear in $\log \|c\|$.

In fact, this algorithm solves (1.1) for any metric μ of the form $d^{H, \ell}$, where $H = (T, U)$ is a sparse frame and ℓ is a nonnegative orbit-invariant function on U . This relies on the following fact.

Statement 5.4 [K97]. *Let m be a minimum 0-extension for G, c, d^H , where H is a frame. Let ℓ be a nonnegative orbit-invariant function on the edges of H , and let*

$\mu = d^{H,\ell}$. Then m is a minimum 0-extension for G, c, μ .

6. Concluding remarks and open questions

Extending a result in [Da] on the intractability of the minimum 3-terminal cut problem (i.e., (1.1) with $\mu = d^{K_3}$), [K98] proves that for a fixed $\mu = d^H$, problem (1.1) is strongly NP-hard if H is non-modular or non-orientable (e.g., if $H = C_5$ or $K_{3,3}$). On the other hand, (1.1) is solvable in strongly polynomial time for each minimizable metric μ . There is one more polynomial case, namely, when μ is representable as $\mu = \lambda_1\mu_1 + \dots + \lambda_k\mu_k$, where $\lambda_1, \dots, \lambda_k \geq 0$, each μ_i is the cut metric on T corresponding to a bi-partition $\{A_i, B_i\}$ of T , and the family $\mathcal{F} = \{A_1, \dots, A_k, B_1, \dots, B_k\}$ satisfies the Helly property, i.e., any subfamily of \mathcal{F} has a nonempty intersection provided that each two members of \mathcal{F} meet (when μ is d^H , this is shown to be equivalent to the property that H is a median graph, i.e., any three nodes of H have precisely one median [MS]). A strongly polynomial algorithm for this case is given in [Ch] (see also [K98, Sec. 5] for a simple algorithm based on a reduction to k minimum cut computations and uncrossing techniques).

Yet, there remain metrics $\mu = d^H$ for which the complexity status of (1.1) is unknown. Let $H = (T, U)$ be modular and orientable but not necessarily hereditary modular, and let Q_1, \dots, Q_k be the orbits of H . Then $\mu = d^H$ is $\mu_1 + \dots + \mu_k$, where μ_i is the corresponding 0-extension of $d^{H/(U-Q_i)}$ to T , and (5.3) holds for any $G = (V, E)$ ($V \supseteq T$) and c , as explained in Section 5. We conjecture that (5.3) holds with equality when each $H/(U - Q_i)$ is a frame with possible parallel edges (a stronger conjecture is that a similar property takes place for any orientable modular graph H).

In this case, each subproblem with G, c, μ_i is solvable in polynomial time, and a method in Section 5 constructs a cyclically even tight extension m' of μ to V satisfying $c \cdot m' \leq \tau_1 + \dots + \tau_k$. If m' is a 0-extension, then m' is an optimal solution to (1.1), and (5.3) turns into equality. However, m' needs not be a 0-extension in general. For example, suppose that H include a cube $H' = (T', U')$ as an isometric subgraph, and let $A \subset T'$ consist of four nodes with the distance two for each pair. Then d^H has a cyclically even tight extension m on $V = T' \cup \{x\}$ such that $m(xv) = 1$ for each $v \in A$ and $m(xv) = 2$ for each $v \in T' - A$ (so m is not a 0-extension in general).

We address an open question: is (1.1) in P when μ is the path metric of an arbitrary orientable modular graph? Two more open questions concern possibility to construct a combinatorial strongly polynomial algorithm for the minimum $(2, r)$ -metric problem, and a combinatorial polynomial algorithm for (1.1) with an arbitrary minimizable μ .

REFERENCES

- [ADK] G.M. Adelson-Velsky, E.A. Dinitz, and A.V. Karzanov, *Flow Algorithms* (Nauka, Moscow, 1975), in Russian.
- [B85] H.-J. Bandelt, Networks with Condorcet solutions, *European J. Oper. Res.* **20** (1985) 314-326.
- [B88] H.-J. Bandelt, Hereditary modular graphs, *Combinatorica* **8** (2) (1988) 149-157.
- [Ch] V. Chepoi, A multifacility location problem on median spaces, *Discrete Applied Math.* **64** (1996) 1-29.
- [Da] E. Dalhaus, D.S. Johnson, C. Papadimitriou, P. Seymour, M. Yannakakis, The complexity of the multiterminal cuts, *SIAM J. Comput.* **23** (4) (1994) 864-894.
- [Dr] A.W.M. Dress, Trees, tight extensions of metric spaces, and the cohomological dimension of certain groups, *Advances in Mathematics* **53** (1984) 321-402.
- [FF] L.R. Ford and D.R. Fulkerson, *Flows in networks* (Princeton Univ. Press, Princeton, NJ, 1962).
- [Is] J. Isbell, Six theorems about metric spaces, *Comment. Math. Helv.* **39** (1964) 65-74.
- [K87] A.V. Karzanov, Half-integral five-terminus flows, *Discrete Applied Math.* **18** (3) (1987) 263-278.
- [K97] A.V. Karzanov, Metrics with finite sets of primitive extensions, *Preprint* 97-110 (SFB 343, Bielefeld University, Bielefeld, 1997) 27p.
- [K98] A.V. Karzanov, Minimum 0-extensions of graph metrics, *European J. Combinatorics* **19** (1998) 71-101.
- [KL] A.V. Karzanov and M.V. Lomonosov, Systems of flows in undirected networks, in: *Mathematical Programming etc.* (Inst. for System Studies, Moscow, 1978, iss. 1) 59-66, in Russian.
- [KM] A.V. Karzanov and Y. Manoussakis, Minimum $(2,r)$ -metrics and integer multiflows, *European J. Combinatorics* **17** (1996) 223-232.
- [Lom] M.V. Lomonosov, Combinatorial approaches to multiflow problems, *Discrete Applied Math.* **11** (1) (1985) 1-94.
- [MS] H.M. Mulder and A. Schrijver, Median graphs and Helly hypergraphs, *Discrete Math.* **25** (1979) 41-50.
- [RW] B. Rothschild and A. Whinston, On two-commodity network flows, *Opera-*

tions Research **14** (1966) 377-387.

[TFL] B.C. Tansel, R.L. Francis, T.J. Lowe, Location on networks: a survey I, II, *Management Sci.* **29** (1983) 482-511.